



The Cloud Feedback Model Intercomparison Project (CFMIP) Progress and future plans

Mark Webb, Keith Williams, Mark Ringer, Alejandro Bodas
Salcedo, Cath Senior (Hadley Centre)

Tomoo Ogura, Yoko Tsushima, (NIES, JAMSTEC Japan)


Sandrine Bony, Majolaine Chiriaco (IPSL/LMD)

and CFMIP contributors (BMRC, GFDL, UIUC, MPI, NCAR)

CERES/GERB meeting October 2006

- Set up by Bryant McAvaney (BMRC), Herve Le Treut (LMD)
- WCRP Working Group on Coupled Modelling (WGCM)
- Systematic intercomparison of cloud feedbacks in GCMs
- +/-2K atmosphere only and 2xCO₂ 'slab' experiments
- Aim to identify key uncertainties
- Link climate feedbacks to cloud observations
- ISCCP simulator required (Klein & Jakob, Webb et al)
- Now have data for 13 GCM versions from 8 groups
- Website shows data available, publications, plans, etc.
- <http://www.cfmip.net>

cfmip homepage - Mozilla Firefox
File Edit View Go Bookmarks Tools Help
http://www.cfmip.net/
Go



Please report any problems with this site to keith.williams@metoffice.gov.uk

<p>Project Overview</p> <p>Detailed Project Description</p> <p>Diagnostic Subprojects</p> <p>Experimental Protocols</p> <p>Data Requirements</p> <p>The ISCCP simulator</p> <p>Project Extensions</p> <p>Participating Groups</p> <p>Data Available</p> <p>FAQ</p> <p>Publications</p> <p>Change History</p>	<p>Lead Co-ordinator: M.J. Webb Hadley Centre, Met Office, UK E-mail: mark.webb@metoffice.gov.uk</p> <p>Co-ordinator: R. Colman Bureau of Meteorology Research Centre (BMRC), Australia E-mail: r.colman@bom.gov.au</p>	<p>Co-ordinator: S. Bony Laboratoire de Météorologie Dynamique (LMD/IPSL), France E-mail: Sandrine.Bony@lmd.jussieu.fr</p> <p>Project Advisor: B.J. McAvaney Bureau of Meteorology Research Centre (BMRC), Australia E-mail: B.McAvaney@bom.gov.au</p>
--	--	---

News

This section contains announcements about the CFMIP project and website. Please email keith.williams@metoffice.gov.uk or mark.webb@metoffice.gov.uk if you wish to add anything.

June 2006 - CFMIP Publications


Work under various CFMIP subprojects is now appearing in the literature. Please see our new publications section. *Mark Webb*

April 2006 - CFMIP Phases I and II

By the time of the fifth IPCC Scientific Assessment we hope that the ISCCP simulator will be required as standard in the IPCC experimental protocol. When this happens the current CFMIP experimental protocol will be redundant. With this in mind we are now thinking about what CFMIP could do in future to develop and apply new techniques for understanding and evaluating cloud climate feedbacks. These ideas will eventually form CFMIP Phase II. It is expected that CFMIP Phase I will continue until the time when daily ISCCP simulator diagnostics are required as standard in the AMIP and CMIP experimental protocols. Further daily data are expected from IPSL, NCAR, and Environment Canada by the end of 2006. *Mark Webb*

Comparison of +/- 2K and slab model experiments

Ringer et al, GRL 2006



Cess experiments capture the spread in cloud feedback from slab experiments.

Offset due to suppressed clear-sky feedbacks in slab vs Cess


Values are global mean changes in NET, SW and LW CRF per degree of warming (Wm⁻²K⁻¹)


Changes in ISCCP cloud types slab vs +/-2K

Ringer et al, 2006




Values are
global mean
change in each
cloud type per
degree of
warming (% / K)






Cloud radiative forcing (CRF) climate response in vertical velocity bins over tropical oceans (30N-30S) from 15 coupled climate models
 8 higher sensitivity and 7 lower sensitivity

Net CRF spread largest in subsidence regions, suggesting low clouds are 'at the heart' of cloud feedback uncertainties



CFMIP cloud feedback “classes”




Models with **positive low cloud feedback** over larger areas have higher sensitivity

Webb et al 2006


Webb et al Climate Dynamics 2006 – CFMIP models



Global Mean




Areas of weak LW cloud feedback




Areas with small **LW** cloud feedbacks explain 59% of the **NET** cloud feedback ensemble variance

Global Mean






Areas of weak LW cloud feedback



Cloud feedbacks in these areas are indeed dominated by reductions in low level cloud amount (shown with ISCCP simulator)

Cloud liquid(2xCO₂ -1xCO₂ kg/kg)

Cloud ice (1xCO₂ kg/kg)



MIROC-HS

MIROC-LS


HadSM4

GFDL

UIUC

Tsushima et al.,2006


Williams et al Climate Dynamics 2006 (CFMIP)



ERBE /ERA SW	ERBE /ERA LW	ERBE/ NCEP SW	ERBE/ NCEP LW	Avg RMS	Avg Climate Sens.& Range
1.2	1.0	1.6	0.9	1.2	3.8K
1.3	1.2	1.3	1.1	1.2	
1.5	1.0	1.9	1.0	1.4	2.9
1.8	1.2	2.1	1.1	1.5	-4.4K
1.7	1.1	2.2	1.2	1.5	
2.1	1.1	2.3	1.0	1.6	3.0K
2.4	1.0	3.0	1.0	1.9	
1.7	1.6	2.2	1.9	1.9	2.3
2.2	1.4	2.8	1.5	2.0	-3.5K
3.4	1.3	3.7	1.1	2.4	

RMS-differences of present-day variability composites against observations for 10 CFMIP/CMIP model versions. The five models with smallest RMS errors tend to have higher climate sensitivities. (Consistent with Bony & Dufresne 2005)

Williams and Tselioudis (Climate Dyn in revision)



ISCCP observational cloud cluster regimes (20N-20S)

In the cloud regime framework, the mean change in cloud radiative forcing can be thought of as having contributions from:

- A change in the RFO (Relative Frequency of Occurrence) of the regime
- A change in the CRF (Cloud Radiative Forcing) within the regime (i.e. a change in the tau-CTP space occupied by the cluster/development of different clusters).

$$\overline{\Delta CRF} = \sum_{i=1}^{nclusters} CRF_i \Delta RFO_i + \sum_{i=1}^{nclusters} RFO_i \Delta CRF_i + \sum_{i=1}^{nclusters} \Delta RFO_i \Delta CRF_i$$

Model	Difference in $\overline{\Delta NCRF}$ (Wm^{-2}/K)	Model λ (Wm^{-2}/K)	Obs. constr. λ (Wm^{-2}/K)	Model clim. Sens. (K)	Obs. constr. Clim. Sens. (K)
ECHAM5	0.49	1.21	0.72	3.3	5.6
HadSM3	0.17	1.06	0.89	3.5	4.2
HadSM4	0.03	1.00	0.97	3.7	3.8
HadGSM1	-0.11	0.83	0.94	4.6	4.1
MIROC-lo	-0.12	0.79	0.91	3.9	3.4
MIROC-hi	-0.19	0.48	0.67	6.5	4.7
Range		0.73	0.30	3.2	2.2
Std. dev.		0.25	0.12	1.2	0.8

CFMIP Phase I continues...



Daily cloud diagnostics to be hosted by PCMDI

To be made available as a community resource

UKMO, MIROC, MPI, NCAR data currently in transit

IPSL, Env Canada promised later this year

Will allow many ISCCP cloud studies to be applied to a representative selection of climate models


Will become part of standard IPCC diagnostic protocol by time of AR5 (agreed by WGCM Sep 06)

CFMIP Phase II – looking further ahead



Co-ordinators: Mark Webb, Sandrine Bony, Rob Colman
Project advisor: Bryant McAvaney

**Main objective : A better assessment of modelled
cloud-climate feedbacks for IPCC AR5**



CFMIP Phase II – planned approach



Develop improved cloud diagnostic techniques in climate models :

- CFMIP CloudSat/CALIPSO simulator
- Cloud water budget / tendency terms
- 3 hourly data at key locations (ARM sites, GPCI)

Explore the sensitivities of cloud feedbacks to differing model assumptions using idealised climate change experiments

Demonstrate the application of these techniques to the understanding and evaluation of cloud climate feedbacks via pilot studies


Organise a systematic cloud feedback model comparison with the next generation of climate models (ideally by embedding suitable cloud diagnostics in the AR5 experimental protocol.)

This is a modular cloud simulator framework which will allow a number of cloud simulator modules to be plugged into climate models via a standard interface.

This is currently under development in collaboration with various groups:

- Hadley Centre (Alejandro Bodas-Salcedo, Mark Webb, Mark Ringer)
- LLNL (Steve Klein, Yuying Zhang)
- LMD/IPSL (Marjolaine Chiriaco, Sandrine Bony)
- CSU (Johnny Lyo, John Haynes, Graeme Stephens)
- PNNL (Roger Marchand)

C3S/CloudSat comparison with UK NWP model (Alejandro Bodas-Salcedo)





Transect through a mature extra-tropical system

Strong signal from ice clouds

Strong signal from precipitation


Cloud and precip not present in obs




ACTSIM LIDAR comparison with GLAS / ICESat data

(M. Chiriaco LMD/IPSL)

Lidar signal **simulated**
from LMDZ outputs



Lidar signal **observed**
from GLAS spatial lidar



→ Evaluation of the vertical structure of the atmosphere in models,
at global scale


Indicates excessive reflectivities from cloud ice in this climate model

Cloud water budget / process diagnostics (Tomoo Ogura NIES Japan)



$$\frac{\partial Q_c(\text{liq} + \text{ice})}{\partial t} = \underbrace{\text{Condensation} + \text{Precipitation}}_{(1)} + \underbrace{\text{Ice-fall-in} + \text{Ice-fall-out}}_{(2)} + \underbrace{\text{Advection}}_{(3)} + \underbrace{\text{Cumulus mixing}}_{(4)} + \underbrace{\text{Conv. adjustment}}_{(5)}$$

- Transient response of terms (1)...(5) to CO₂ doubling is monitored every year.
- Terms showing positive correlation with cloud water response contribute to the cloud water variation.




Qc increase related to the source term

Qc decrease related to the source term

Cloud water budget diagnosis in MIROC (Tomoo Ogura NIES Japan)



Correlation coefficient (when positive)
[Cloud water vs (1)Condens-Precip]




Decrease in mid-low level sub-tropical cloud water correlates with decrease in large-scale condensation


GCSS/WGNE Pacific Cross-section Intercomparison (GPCI)

Joao Teixeira teixeira@nurc.nato.int

ISCCP total cloud cover



Sea Surface Temperature



GPCI is a working group of the GEWEX Cloud System Study (GCSS)


Models and data are analyzed along a Pacific Cross section from Stratocumulus, to Cumulus and to deep convection

Period: June-July-August 1998 and 2003

Time resolution: 0, 3, 6, 9, 12, 15, 18, 21 UTC

Mean GPCI liquid water cross section - JJA98

from three climate models



Too shallow -> fog

Is this too much
liquid water?


How deep should
the PBL be..?

Joao Teixeira teixeira@nurc.nato.int

Mean diurnal cycle: ISCCP cloud cover

Joao Teixeira teixeira@nurc.nato.int

peak values of Sc cloud
cover around 32-35 N




Diurnal cycle: max in
(early) morning local time

peak values of mid/high
clouds close to ITCZ

JJA98 mean diurnal cycle: model low cloud cover

Joao Teixeira teixeira@nurc.nato.int



peak values too far to
the south (around 26 N)

realistic diurnal
cycle: morning max

Proposed point diagnostics for CFMIP Phase II



- 3 hourly instantaneous model data
- 10-20 years of data to give stable statistics
- on a grid of locations covering the GPCI and TWP
- additional key locations (e.g. ARM sites, high latitudes)
- in AMIP and idealised climate change experiments

The aim is to align climate model diagnostics with those in use in GCSS/ARM to encourage a wider group of people to examine and criticise cloud feedbacks in climate models

For example this would give insight into the impact of diurnal cycle errors on cloud-climate feedbacks

Sensitivity tests are proposed using the idealised climate change experiments developed by Brian Soden:

i.e. AMIP and AMIP + composite CMIP SST anomaly

The aim is to quantify the impact of certain differences in model formulation on cloud-climate feedbacks by implementing consistent treatments across models

Possible examples include:

- fixed liquid cloud water content and radiative properties
- consistent precipitation & mixed phase partitioning
- consistent boundary layer resolution
- consistent simple shallow convection scheme?

Concrete project proposal by Jan 2007

Aim for endorsement by WGCM and GEWEX SSG
in early 2007

Joint CFMIP/ENSEMBLES meeting Paris April 2007

Development of diagnostics / pilot studies 2007-2008

Systematic model inter-comparison with new model
versions 2008- (preferably as part of AR5 models)

CFMIP Phase I – mostly ISCCP/ERBE/ISCCP_FD

All of the following will benefit CFMIP Phase II:

- CERES SRBAVG GEO, SYN/AVG/ZAVG products

- CERES/MODIS cloud retrievals


- CloudSat/CALIPSO/CERES merged products

- Surface + ATM + TOA products (e.g. SARB)

- Diurnal cycle GEO/GERB data

The barriers are often in bringing the sampling and statistical summaries from satellite products and GCM diagnostics into line – e.g. providing ISCCP D1-like tau-Pc histograms


Hawaii – California transect: March 2005



mark.webb@metoffice.gov.uk

Cleaner separation of SW feedbacks

Taylor et al (in revision)




... APRP (Approximate Partial Radiative Perturbation) method for separation of SW cloud and non-cloud feedbacks.

... Validated against full GFDL PRP calculations courtesy of B. Soden.

Similar to Yokohata et al method but with subtle differences.

Fig. 1: Schematic representation of a simple shortwave radiation model showing fluxes passing through the atmosphere and being partially reflected on each pass, where S is the insolation, α surface albedo, and γ and μ are the atmospheric scattering coefficient and transmissivity, respectively. Wavelengths that are readily absorbed by the atmosphere are assumed to be completely removed on the first pass, and the atmosphere is transparent to other wavelengths.

Taylor et al SW APRP on CFMIP model data



APRP method gives a more positive SW cloud feedback and larger spread than CRF method

This is because it makes a cleaner separation between surface albedo and cloud feedbacks

Michel Crucifix, Karl Taylor, Mark Webb

Summary

















A number of studies now point to low cloud feedbacks being a key uncertainty in climate models

Daily cloud/radiation/ISCCP simulator diagnostics from CFMIP Phase I will be available from PCMDI by the end of the year



Reductions in uncertainty due to cloud feedbacks will require links to observations and also understanding and criticism of feedback mechanisms in models

CFMIP Phase II is an opportunity to align diagnostics in a range of models with those in use in GCSS/ARM/CPT


We hope to formalise our plans by the end of this year.




SW CRF response along Pacific transect in CFMIP and AR4 slab models. Cloud feedbacks are typically smaller than model biases. No clear relationship between bias and feedback




Large-Scale Cloud Fraction HadSM3 Control [149W,17N]




Large-Scale Cloud Fraction HadSM3 2CO2 [149W,17N]




Convective Cloud Fraction HadSM3 Control [149W,17N]



Convective Cloud Fraction HadSM3 2CO2 [149W,17N]







Interannual sensitivity of CRF to SST in vertical velocity bins over tropical oceans (30N-30S) from 15 coupled climate models
8 higher sensitivity and 7 lower sensitivity and observed (ISCCP FD)

Models underpredict low cloud sensitivity, but higher sensitivity models less so.



Low cloud response in HadSM3 transition regions



- Deep and shallow convection weaken in the warmer climate (consistent with Held and Soden 2006)
- Shallow convection typically detrains into two model layers in present day, but one level in the warmer climate
- If a certain amount of water vapour is detrained into a single layer it will moisten that layer more than would be the case if it was spread over two layers
- May explain why HadSM3 stratiform cloud fraction increases with weakening shallow convection
- Hence the negative low cloud feedback in HadSM3 may be due to poor vertical resolution and so not credible





Large-Scale Cloud Fraction
HadGSM1 Control [149W,17N]




© Crown copyright 2006

Large-Scale Cloud Fraction
HadGSM1 2CO2 [149W,17N]




Page 37



Requires: more cloud diagnostics from GCMs + enhanced scrutiny

CFMIP-II:

- Encourage the analysis of cloud feedback processes by a wider community !
 - make cloud diagnostics more easily accessible to the community (daily cloud diagnostics from CFMIP-1 to be available from PCMDI)
 - strengthen the link between CMIP and CFMIP (e.g. by increasing the number of cloud diagnostics in the outputs of coupled models, by running the ISCCP simulator)
- Organize climate physics sensitivity experiments + consistent implementation of simplified physics (e.g. mixed-phase cloud feedback)



How may we use our *physical understanding* of climate change cloud feedbacks and the available *model-data comparisons* to define a “metrics” for cloud feedbacks?

CFMIP-II:

- Explore relationships between cloud evaluation tests and cloud-climate feedbacks based on a wide diversity of diagnostics and approaches + a large number of GCMs
- Discuss the issue during a joint [CFMIP/ENSEMBLES workshop in Paris \(11-13 April 2007\)](#).

CFMIP Phase II will aim to reduce uncertainty in cloud feedbacks and climate sensitivity by developing further links between feedbacks and observations **and** improving our understanding of cloud feedback mechanisms by:

1. Developing better cloud diagnostics for models:
 - CloudSat/CALIPSO simulator
 - GCSS Pacific Cross Section (diurnal cycle, ...)
 - Cloud budget/tendency diagnostics
2. Applying
2. Exploring sensitivities of feedbacks to model physics
e.g. low clouds, convective entrainment
3. Collaboration with Gewex/GCSS community
 - GCSS Pacific Cross Section (diurnal cycle, ...)

Cess +/-2K fixed season experiments are not a quantitative guide to coupled model feedbacks

- no seasonal cycle
- no high latitude amplification of warming

Alternative options include:

- continuing use of mixed layer experiments
- re-running sections of AR4 coupled experiments with extra diagnostics
- 'patterned SST perturbation' experiments as developed by Brian Soden (possibly AMIP+)

CFMIP Cloudsat/CALIPSO Simulator (C3S)

Sub-timestep information


- Cloud condensate budget terms
- Physics increments for temperature, humidity, etc

Detailed diagnostics at key locations as used in
GCSS/ARM studies (e.g GPCI)

European Cloud System study (EUROCS) Pacific Cross Section



A. P. SIEBESMA *et al.*




Siebesma et al 2004


Pacific Transect

California/trades/ITCZ

9 regional, NWP and climate
models compared, JJA 1998

© Crown copyright 2006






Develop better cloud diagnostics for climate models:

High frequency instantaneous diagnostics along
WGNE/GCSS GPCI and at ARM sites

CFMIP CloudSat/CALIPSO Simulator (C3S)



Requires: process-level studies + model-data comparisons + new satellite data

→ [Embed GCSS](#) (e.g. high-frequency diagnostics at ARM sites, GEWEX Pacific Cross-Section; evaluation of some specific cloud processes)

→ [Development of a CFMIP CloudSat/CALIPSO Simulator](#) (C3S) (including eventually an A-train orbital simulator) -> will favor interactions with obs people!

→ We would like the [AR5 models](#) to be run with the [ISCCP/radar/lidar simulator](#), and the [CFMIP cloud diagnostics](#) to be included into the list of standard outputs

Reducing uncertainty by understanding feedbacks



- 1/ Try to understand how physical cloud climate feedback processes are operating in models
- 2/ Ask 'Is this behaviour credible?'
- 3/ Develop new schemes with more credible cloud-climate feedback behaviour in mind
- 4/ Differences between model feedbacks may reduce in the longer term
- 5/ Can help to focus attention on key physical processes for cloud feedbacks

New diagnostics will provide opportunities for new links

e.g. CloudSat/CALIPSO data may constrain models with strong mixed-phase cloud feedbacks due to excessive amounts of cloud ice

This will be an ongoing area of research, and the focus of a joint ENSEMBLES/CFMIP meeting in Paris 11th-13th April 2007

CFMIP CloudSat/CALIPSO simulator (Alejandro Bodas-Salcedo)




Simulated CloudSat reflectivities from UKMO forecast model

The simulator consists of 5 steps:

1. Orbital simulation
2. Sampling
3. Preprocessing
4. Subgrid sampling of cloud overlaps
5. Radar reflectivity calculated using code provided by Matt Rogers (CSU)





Outputs:

- Reflectivity from clouds and precipitation (without attenuation)
- Total reflectivity, accounting for attenuation by gases, clouds and precipitation
- Products obtained from inputs at gridbox and subgrid scales



ACTSIM CALIPSO simulator

— Marjolaine Chiriaco (LMD/IPSL)



Initially we will provide an A-train orbital simulator to allow climate modellers to save model cloud variables co-located with CloudSat/CALIPSO overpasses

Initially sampled data would be submitted to CFMIP and both simulators run centrally

As the approach matures we plan to integrate this package with the ISCCP simulator so that it can be run in-line as part of the model development cycle